

## **ISOLATION, CHARACTERIZATION AND ENUMERATION OF EFFICIENT MICROORGANISMS IN TERMITE MOUND MATERIALS**

**Dr.P.SATHIYA BAMA,**

Associate Professor and Head, Department of Zoology,

M.V.Muthiah Government Arts College for Women, Dindigul (Dt), Tamilnadu, India.

E-mail : yaeshabna@rediffmail.com

### **ABSTRACT**

*This work highlights the isolation, identification and characterization of beneficial microbial population in ten different termite mound materials. As major eco-engineers in tropical ecosystems, termites build biogenic structures with galleries, sheeting's, fungus-comb chambers where the diazotrophs influence nitrogen fixation altering soil mineralogy. Nitrogen fixers strongly influence the physical and chemical properties of soils there by SOM gets altered. In dry land ecosystems, termite mound materials (TMM) are used to enhance soil fertility thus mound soil acts as hotspots for primary production.*

### **INTRODUCTION**

Termites are the dominant arthropod decomposers in tropical ecosystems. Termites are major components of detritivore macrofauna feeding on a whole range of living, recently dead plant material in various stages of decomposition (Wood, 1988). Among the soil invertebrates, termites act as ecosystem engineers (Jones *et al.* 1994) which have profound effect on promotion of litter decomposition and the formation of stable pools of soil organic matter.

Termites perform several activities that qualify them as soil engineers. Free living N<sub>2</sub>-fixing bacteria are found to be associated in termite gut and mound soil. Biological nitrogen fixation offers a non polluting source of nitrogen and can improve crop production and decrease the global use of synthetic fertilizers. Besides this, termite mound material (TMM) can be used as organic resources to improve soil productivity. It can also counteract land degradation through their soil borrowing and feeding activities.

### **MATERIALS AND METHODS**

Isolation enumeration and characterization of indigenous isolates of *Azotobacter*, *Beijerinckia*, *Derxia* sp, Actinomycetes and heterotrophs from different termite mound material

VISIT...

LANZAROTE  
*Caliente*.COM

(*Trinervius trinervoides* and *Odontotermes obes*) in the vicinity of various blocks of Dindigul district, TamilNadu, India were carried out. The population of the different groups of microorganisms such as bacteria and actinomycetes in various samples were enumerated using standard plate count method (Kannan and Rao, 1996).

One gram of material from each termite mound soil were suspended in 99 ml of sterile distilled water in a 250 ml Erlenmeyer flask and shaken vigorously for 30 minutes with orbital shaker incubator at 150 rev / min at 24°C to form a uniform solution of  $10^{-2}$  concentration and served as master stock. This stock was subjected to serial dilution with sterile distilled water using a sterile pipette to form  $10^{-3}$  to  $10^{-7}$  concentration. One ml of dilutions  $10^{-3}$  /  $10^{-4}$  were pipetted out to sterile petri plate for the enumeration of bacterial count; the same was repeated with dilutions of  $10^{-5}$  /  $10^{-6}$  for actinomycetes. The media and growth condition used for the enumeration of Total Colony Forming Units (CFU/ml) of bacteria and actinomycetes are given as follows

**Table 1: Media and growth condition**

| Sl. No. | Microbial group      | Isolation media          | Dilution factor   | Growth temperature (°C) | Period of incubation (in days) |
|---------|----------------------|--------------------------|-------------------|-------------------------|--------------------------------|
| 1       | <i>Azotobacter</i>   | Waksman                  | $10^{-3}/10^{-4}$ | 37°C                    | 7                              |
| 2       | <i>Beijerinckia</i>  | Jensens                  | $10^{-3}/10^{-4}$ | 37°C                    | 5                              |
| 3       | <i>Derxia</i>        | Cample's and Doberneir's | $10^{-3}/10^{-4}$ | 37°C                    | 5                              |
| 4       | <i>Actinomycetes</i> | Kenknight's Agar         | $10^{-3}/10^{-4}$ | 37°C                    | 5-7                            |
| 5       | Heterotrophs         | Soil extract agar        | $10^{-5}/10^{-6}$ | 37°C                    | 2-3 days                       |

After incubation counts of each microbial groups were noted, average counts and the total population per ml of sample were calculated. The microbial population was expressed as Colony Forming Units (CFU / ml) of the sample. The distinct viable colonies were picked and restreaked on to appropriate agar medium to obtain pure cultures. Each isolate showed its characteristic growth, pigmentation and biochemical reactions. All the bacterial and actinomycetes isolates were identified through morphological and biochemical characteristics. The parameters investigated include colony morphology, gram's reactions, motility, acid production, methyl red reaction, green fluorescent. Voges Proskaur (VP) reaction, catalase reaction, cellulose activity and starch hydrolysis. (Apun., 2000). The test results were compared with that of Bergey's Manual of Determinative Systematic Bacteriology (Holt et.al, 1987) for identification of bacterial and actinomycetes isolates.

## RESULTS AND DISCUSSIONS

The isolation, characterization and enumeration of efficient microbes like diazotrophs from termite mound material (TMM) of different species are given in table no. 2 and 3. The total colony forming units of diazotrophs varied across the different samples.

**Table 2 : Characterisation of diazotrophs isolated from termite mound material**

| Colony characters.        | <i>Azotobacter</i> sp                                            | <i>Beijerinckia</i> sp                                                             | <i>Derxia</i> sp                           |
|---------------------------|------------------------------------------------------------------|------------------------------------------------------------------------------------|--------------------------------------------|
| Cell shape                | Ovoid ,rod or cocci appears in pairs                             | Straight rods with rounded ends                                                    | Rods                                       |
| Colony character          | Smooth opaque, convex circular gummy colony with undulate margin | Smooth irregular folded glistening and raised colony with tenacious elastic slime. | White to brown. extremely tenacious growth |
| Cell size                 | 2 x 1.5µm                                                        | 2x1 µm                                                                             | 1.0-1.2 µm                                 |
| Gram reaction             | G-ve                                                             | G-ve                                                                               | G-ve                                       |
| Flagella                  | Peritrichous                                                     | Peritrichous                                                                       | Peritrichous                               |
| Motility                  | +                                                                | +                                                                                  | +                                          |
| Green fluorescent         | +                                                                | +                                                                                  | -                                          |
| Acid production           | +                                                                | +                                                                                  | +                                          |
| Cellulose hydrolysis      | -                                                                | -                                                                                  | -                                          |
| Starch hydrolysis         | +                                                                | +                                                                                  | -                                          |
| Glucose utilization       | +                                                                | +                                                                                  | -                                          |
| Sucrose utilization       | +                                                                | +                                                                                  | -                                          |
| Maltose utilization       | -                                                                | -                                                                                  | -                                          |
| Biotin requirement        | -                                                                | -                                                                                  | -                                          |
| Cyst formation            | +                                                                | +                                                                                  | -                                          |
| Polysaccharide production | +                                                                | +                                                                                  | +                                          |

The population of *Azotobacter* sp registered a maximum in TM4, TM5 and TM3 of 8.37, 8.17 and 8.15 ( $\times 10^3$  CFU/g) respectively. The population of *Beijerinckia* registered a maximum of 5.67, 5.2 and 5.13 ( $\times 10^3$  CFU/g) in TM4, TM3 and TM5 material. The *Derxia* load ( $\times 10^3$  CFU / g) varied from 1.7 to 2.87 in different termite mound analysed. Compared to *Azotobacter* and *Beijerinckia*, *Derxia* load was lesser than other diazotrophs. Highest number of *Azotobacter* and *Beijerinckia* population were recorded in TM5 and TM4. Likewise, the actinomycetes population

registered a maximum load of 12.38 ,12.23 and 12.12 ( $\times 10^3$  CFU / g) in TM5, TM3 and TM4 respectively. The heterotrophs population were maximum in TM5 and TM3 with 41.67 and 40.27 ( $\times 10^6$  CFU/g). On analysis, the TM4, TM3 and TM5 were found to possess a maximum diazotroph and microbial load than other mound samples.

**Table 3 : Enumeration of microbes isolated from different termite mound at various selected sites**

| Termite mound | <i>Azotobacter</i> sp $\times 10^3$ | <i>Beijerinckia</i> sp $\times 10^3$ | <i>Derxia</i> sp $\times 10^3$ | Actinomycetes $\times 10^3$ | Heterotrophs $\times 10^6$ |
|---------------|-------------------------------------|--------------------------------------|--------------------------------|-----------------------------|----------------------------|
| TM1           | 5.10<br>$\pm 0.20$                  | 4.37<br>$\pm 0.12$                   | 1.23<br>$\pm 0.17$             | 9.50<br>$\pm 0.01$          | 26.77<br>$\pm 0.14$        |
| TM2           | 6.43<br>$\pm 0.12$                  | 3.60<br>$\pm 0.36$                   | 1.37<br>$\pm 0.12$             | 9.45<br>$\pm 0.02$          | 29.77<br>$\pm 0.12$        |
| TM3           | 8.15<br>$\pm 0.12$                  | 5.20<br>$\pm 0.08$                   | 2.57<br>$\pm 0.12$             | 12.23<br>$\pm 0.02$         | 40.27<br>$\pm 0.12$        |
| TM4           | 8.37<br>$\pm 0.12$                  | 5.67<br>$\pm 0.12$                   | 2.87<br>$\pm 0.12$             | 12.12<br>$\pm 0.07$         | 38.07<br>$\pm 0.05$        |
| TM 5          | 8.17<br>$\pm 0.12$                  | 5.13<br>$\pm 0.12$                   | 2.50<br>$\pm 0.08$             | 12.38<br>$\pm 0.04$         | 41.67<br>$\pm 0.12$        |
| TM 6          | 4.63<br>$\pm 0.17$                  | 2.17<br>$\pm 0.12$                   | 1.35<br>$\pm 0.01$             | 10.17<br>$\pm 0.15$         | 34.17<br>$\pm 0.24$        |
| TM 7          | 6.57<br>$\pm 0.12$                  | 4.43<br>$\pm 0.12$                   | 1.17<br>$\pm 0.12$             | 8.86<br>$\pm 0.01$          | 27.57<br>$\pm 0.34$        |
| TM 8          | 5.47<br>$\pm 0.12$                  | ND                                   | 1.47<br>$\pm 0.12$             | 8.23<br>$\pm 0.01$          | 26.17<br>$\pm 0.12$        |
| TM 9          | 5.47<br>$\pm 0.12$                  | 1.80<br>$\pm 0.08$                   | ND                             | 8.77<br>$\pm 0.12$          | 24.30<br>$\pm 0.08$        |
| TM 10         | 4.23<br>$\pm 0.12$                  | 2.67<br>$\pm 0.12$                   | 1.77<br>$\pm 0.12$             | 8.10<br>$\pm 0.01$          | 25.60<br>$\pm 0.36$        |

ND - Not detected Values are mean  $\pm$  standard error (n = 3)

Microorganisms are the dominant biotic structural components in soil and have higher biomass specific activities. The list of N<sub>2</sub> fixing bacteria associated with non legumes includes species of *chromobacter*, *Acetobacter*, *Azotobacters*, *Beijerinckia*, *Bacillus*, *Enterobacter*, *Erwinia*, *Derxia* and *Rhodospirillum* (Wani,1990). Although many genera and species of N<sub>2</sub> fixing bacteria are isolated from the rhizosphere of various cereals, mainly members of *Azotobacter* and *Azospirillum* genera have been widely used to increase the yield of cereals and legumes in green house and under field conditions.

Diazotrophs are found in a wide variety of habitats including free living forms in soils, water, termite mound material (TMM), cyanobacterial symbioses with various plants and root nodule symbiosis with legumes (Dixon and Kahn, 2004). *Azotobacter* is a free living nitrogen fixing bacterium, which is used as a biofertilizer in the cultivation of most crops. It has several metabolic capabilities with highest metabolic rate to fix 20-60 kg nitrogen per hectare of land annually. *Azotobacter* is the most common biofertilizer for plants like maize, wheat, sorghum and rice which produces some plant growth promoting metabolites, nutrients to plants, enzymes, different growth hormones like IAA, auxins, gibberellins, cytokinins, vitamins and siderophores. It is probable that non symbiotic nitrogen fixers of the microbial population are widespread and abundant in termite mound material (TMM) which accounts for higher nitrogen fixation and crop yield.

## CONCLUSION

Non symbiotic diazotrophs in termite mound can promote economic and environmental benefits including increased income from high yield, reduce fertilizer cost and reduce leaching of  $\text{NO}_3$  to ground water. In this regard, diazotrophic diversity of termite mound material (TMM) and characterization are keys to understand the significance of nitrogen fixers as biofertiliser. Various environmental determinants and factors inherent to termite biology influence microbial distribution and play a role in ecosystem processing influencing soil fertility by providing beneficial role in agriculture.

## REFERENCES

- Apun, K., Jong, B.C., and Salleh, M.A. 2000 .Screening and isolation of a cellulolytic and amylolytic *Bacillus* from sago pith waste. *J. Gen. Appl. Microbial.* 46 : 263 – 267.
- Dixon,R.and Kahn,D.,2004.Genetic regulation of biological nitrogen fixation. *Nat Rev Microbiol.*2:621–31.
- Holt,J.,A. 1987. Carbon mineralization in semi arid northeastern Australia: The role of termites. *J. Trop. Ecol.* 3:255-263.
- Jones,C,G.,Lawton, J. H. and Shaghak, M.,1994 .Organisms as ecosystem engineers. *Oikos*, 69, 373–386.
- Kannan, N., and Rao .1996. ‘Laboratory Manual in General Microbiology’, Palani Paramount Publication, India.
- Wani S.P.1990.Inoculation with associative nitrogen fixing bacteria: Role in cereal grain production Improvement. *Indian J. Microbiol.* 30: 363-393.
- Wood, T. 1988. Termites and the soil environment. *Biol. Fertility soils.* 6:228 –236.